

Improving Great Lakes Water Balance Forecasting through Regionalized Calibration of the Large Basin Runoff Model

Lauren Fry and Andrew Gronewold

lauren.fry@noaa.gov

drew.gronewold@noaa.gov

Cooperative Institute for Limnology and Ecosystems Research, University of
Michigan

Great Lakes Environmental Research Laboratory, National Oceanic and
Atmospheric Administration

Great Lakes Operational Meteorology Workshop
Chicago, IL
March 14-16, 2012

Outline

1 Introduction (Runoff and the Great Lakes Water Balance)

Outline

- 1 Introduction (Runoff and the Great Lakes Water Balance)
- 2 Improving the Large Basin Runoff Model

Outline

- 1 Introduction (Runoff and the Great Lakes Water Balance)
- 2 Improving the Large Basin Runoff Model
- 3 Summary

Outline

- 1 Introduction (Runoff and the Great Lakes Water Balance)
- 2 Improving the Large Basin Runoff Model
- 3 Summary

WATERBORNE COMMERCE


A 3D coordinate system with three axes. The vertical axis is labeled 'Upbound' at the top and 'Downbound' at the bottom. The horizontal axis is labeled 'Downbound' at the right. The third axis, pointing diagonally down and to the left, is unlabeled. The origin is marked with '0'. Two points are marked on the vertical axis: '40 000 000' and '20 000 000'.






Predicting Water Levels - Great Lakes AHPS

Great Lakes Advanced Hydrologic Prediction System (AHPS)




Predicting Water Levels - Great Lakes AHPS


Great Lakes Advanced Hydrologic Prediction System (AHPS)




Large Basin Runoff Model




Observing Runoff




Observing Runoff




Observing Runoff




Observing Runoff




Observing Runoff




Observing Runoff



Observing Runoff




Observed Runoff Estimates (Area Ratio Method)



$$Q_{subbasin} = \frac{Area_{subbasin}}{\sum Area_{gauge}} * \sum Q_{gauge}$$

Evaluation of Previous LBRM Calibration



Outline

- 1 Introduction (Runoff and the Great Lakes Water Balance)
- 2 Improving the Large Basin Runoff Model
- 3 Summary

It's time to recalibrate!

It's time to recalibrate!

- New ways of estimating parameters


It's time to recalibrate!

- New ways of estimating parameters
- New (more, and more recent) data for calibration


It's time to recalibrate!

- New ways of estimating parameters
- New (more, and more recent) data for calibration
- New ways to incorporate geospatial information on drivers of hydrologic response

Ungauged Basins



Ungauged Basins



What drives hydrologic response?

What drives hydrologic response?

- Hydromorphology

What drives hydrologic response?

- Hydromorphology
 - Stream Density
 - Basin Area
 - Basin Compactness
 - Strahler Stream Order

What drives hydrologic response?

- Hydromorphology
 - Stream Density
 - Basin Area
 - Basin Compactness
 - Strahler Stream Order
- Subsurface Properties

What drives hydrologic response?

- Hydromorphology
 - Stream Density
 - Basin Area
 - Basin Compactness
 - Strahler Stream Order
- Subsurface Properties
 - Soil Thickness
 - Soil Hydraulic Properties
 - Basin Geology

What drives hydrologic response?

- Hydromorphology
 - Stream Density
 - Basin Area
 - Basin Compactness
 - Strahler Stream Order
- Subsurface Properties
 - Soil Thickness
 - Soil Hydraulic Properties
 - Basin Geology
- Land Cover

What drives hydrologic response?

- Hydromorphology
 - Stream Density
 - Basin Area
 - Basin Compactness
 - Strahler Stream Order
- Subsurface Properties
 - Soil Thickness
 - Soil Hydraulic Properties
 - Basin Geology
- Land Cover
 - Percent Forest
 - Percent Agriculture
 - Percent Developed

What drives hydrologic response?

- Hydromorphology
 - Stream Density
 - Basin Area
 - Basin Compactness
 - Strahler Stream Order
- Subsurface Properties
 - Soil Thickness
 - Soil Hydraulic Properties
 - Basin Geology
- Land Cover
 - Percent Forest
 - Percent Agriculture
 - Percent Developed
- Climate

What drives hydrologic response?

- Hydromorphology
 - Stream Density
 - Basin Area
 - Basin Compactness
 - Strahler Stream Order
- Subsurface Properties
 - Soil Thickness
 - Soil Hydraulic Properties
 - Basin Geology
- Land Cover
 - Percent Forest
 - Percent Agriculture
 - Percent Developed
- Climate
 - Precipitation (quantity, seasonality, percent as snow)
 - Temperature

New Methods for Prediction in Ungauged Basins

New Methods for Prediction in Ungauged Basins

- How does hydrologic response vary across the basin?


New Methods for Prediction in Ungauged Basins

- How does hydrologic response vary across the basin?
- Can we use relationships between hydrologic response and watershed characteristics to predict runoff in ungauged portions?


New Methods for Prediction in Ungauged Basins

- How does hydrologic response vary across the basin?
- Can we use relationships between hydrologic response and watershed characteristics to predict runoff in ungauged portions?
- New PUB methods calibrate by constraining model output to hydrologic response based on watershed characteristics


Hydrologic Response



Hydrologic Response




Hydrologic Response




Hydromorphology

Hydromorphology

Q to P vs. Stream Density




Hydromorphology




Subsurface Properties

Subsurface Properties

Q to P vs. Depth to Bedrock




Subsurface Properties




Land Cover

Land Cover

Q to P vs. Percent Agriculture




Land Cover




Climate

Climate

Q to P vs. Average Annual Precipitation




Climate




Climate

Q to P vs. Percent Precipitation as Snow



Climate



Future Work

Future Work

- 1 Determine relationships between hydrologic response and catchment characteristics at gauged watersheds

Future Work

- 1 Determine relationships between hydrologic response and catchment characteristics at gauged watersheds
- 2 Recalibration at three test gauges using PUB methodology and new parameter estimation methods: Clinton, Saginaw, Grand (MI)

Future Work

- 1 Determine relationships between hydrologic response and catchment characteristics at gauged watersheds
- 2 Recalibration at three test gauges using PUB methodology and new parameter estimation methods: Clinton, Saginaw, Grand (MI)
- 3 Expand to entire Great Lakes Basin

Future Work

- 1 Determine relationships between hydrologic response and catchment characteristics at gauged watersheds
- 2 Recalibration at three test gauges using PUB methodology and new parameter estimation methods: Clinton, Saginaw, Grand (MI)
- 3 Expand to entire Great Lakes Basin
- 4 Compare with other models?

Outline

- 1 Introduction (Runoff and the Great Lakes Water Balance)
- 2 Improving the Large Basin Runoff Model
- 3 Summary

Summary

- LBRM is used to provide runoff estimates to the GL AHPS

Summary

- LBRM is used to provide runoff estimates to the GL AHPS
- Calibration of LBRM can be improved by incorporating catchment characteristics

Summary

- LBRM is used to provide runoff estimates to the GL AHPS
- Calibration of LBRM can be improved by incorporating catchment characteristics
- Future work will provide a new and improved calibration that includes estimates of uncertainty

Acknowledgements

- Tim Hunter, Alicia Ritzenthaler
- Cooperative Institute for Limnology and Ecosystems Research
- NOAA Great Lakes Environmental Research Laboratory

Improving Great Lakes Water Balance Forecasting through Regionalized Calibration of the Large Basin Runoff Model

Lauren Fry and Andrew Gronewold

lauren.fry@noaa.gov

drew.gronewold@noaa.gov

Cooperative Institute for Limnology and Ecosystems Research, University of
Michigan

Great Lakes Environmental Research Laboratory, National Oceanic and
Atmospheric Administration

Great Lakes Operational Meteorology Workshop
Chicago, IL
March 14-16, 2012